

Effects of Using a Recurrent Adverse Drug Reaction Prevention Program on Safe Use of Medicine among Patients Receiving Services at the Accident and Emergency Department of Songkhla Hospital Thailand

Authors : Thippharat Wongsilarat, Parichat tuntilanon, Chonlakan Prataksitorn

Abstract : Recurrent adverse drug reactions are harmful to patients with mild to fatal illnesses, and affect not only patients but also their relatives, and organizations. To compare safe use of medicine among patients before and after using the recurrent adverse drug reaction prevention program . Quasi-experimental research with the target population of 598 patients with drug allergy history. Data were collected through an observation form tested for its validity by three experts (IOC = 0.87), and analyzed with a descriptive statistic (percentage). The research was conducted jointly with a multidisciplinary team to analyze and determine the weak points and strong points in the recurrent adverse drug reaction prevention system during the past three years, and 546, 329, and 498 incidences, respectively, were found. Of these, 379, 279, and 302 incidences, or 69.4; 84.80; and 60.64 percent of the patients with drug allergy history, respectively, were found to have caused by incomplete warning system. In addition, differences in practice in caring for patients with drug allergy history were found that did not cover all the steps of the patient care process, especially a lack of repeated checking, and a lack of communication between the multidisciplinary team members. Therefore, the recurrent adverse drug reaction prevention program was developed with complete warning points in the information technology system, the repeated checking step, and communication among related multidisciplinary team members starting from the hospital identity card room, patient history recording officers, nurses, physicians who prescribe the drugs, and pharmacists. Including in the system were surveillance, nursing, recording, and linking the data to referring units. There were also training concerning adverse drug reactions by pharmacists, monthly meetings to explain the process to practice personnel, creating safety culture, random checking of practice, motivational encouragement, supervising, controlling, following up, and evaluating the practice. The rate of prescribing drugs to which patients were allergic per 1,000 prescriptions was 0.08, and the incidence rate of recurrent drug reaction per 1,000 prescriptions was 0. Surveillance of recurrent adverse drug reactions covering all service providing points can ensure safe use of medicine for patients.

Keywords : recurrent drug, adverse reaction, safety, use of medicine

Conference Title : ICNM 2015 : International Conference on Nursing Management

Conference Location : London, United Kingdom

Conference Dates : May 25-26, 2015