

Organic Geochemical Characterization of the Ordovician Source Rock in the Chotts Basin, Southern Tunisia

Authors : Anis Belhaj Mohamed, Moncef Saidi, Mohamed Soussi, Ibrahim Bouazizi, Monia Ben Jrad

Abstract : This paper summarizes the results of Rock-Eval pyrolysis and biomarker data of shale samples collected from the Ordovician age (Llanvirnian-Llandeilian) (Azzel Formation) in the Chotts basin southern part of Tunisia. The results are supported by analysis of cutting samples from wells. The Azzel shales has poor to moderate, occasionally good, potential for sourcing oil and gas with Total Organic Carbon (TOC) content varying from 0.80 to 4.49 % and petroleum potential (PP) values varying between 0.68 to 9.20 Kg of HC/t rock in Baguel and Alaguia wells. However, the Azzel Formation show poor to fair TOC and PP in Elfranig and HajBrahim wells not exceeding 1.10% and 1.05 kg HC/t of rock respectively. The Hydrogen Index (HI) and the Oxygen Index (OI) values of 95-165 mg S₂/g TOC and of 33-108 mg CO₂/g rock relatively show that the Ordovician shales exhibit type II Kerogen that reached the main oil window stage and that the organic matter was bad preserved, Tmax values of 435 - 448°C indicate the organic matter is mature. The biomarker features of the extract samples are characterized by high proportion of tricyclic terpanes that are dominated by C₂₃ and C₂₁ tricyclic terpanes. The hopanes fraction is dominated by C₂₉ and C₃₀ hopanes. The Ordovician shales show a predominance of C₂₇ over C₂₉ steranes (C₂₇/C₂₉>1) and relatively high proportions of diasteranes supporting the shaly character of the source rock.

Keywords : biomarkers, organic geochemistry, ordovician source rock, diasteranes

Conference Title : ICSRD 2020 : International Conference on Scientific Research and Development

Conference Location : Chicago, United States

Conference Dates : December 12-13, 2020