

Effect of Palm Oil Mill Effluent on Microbial Composition in Soil Samples in Isiala Mbano Lga

Authors : Eze Catherine Chinwe, J. D. Njoku

Abstract : Background: Palm oil mill effluent is the voluminous liquid waste that comes from the sterilization and clarification sections of the oil palm milling process. The raw effluent contains 90-95% water and includes residual oil, soil particles, and suspended solids. Palm oil mill effluent is a highly polluting material and much research has been dedicated to means of alleviating its threat to the environment. Objectives: 1. To compare Physico-chemical and microbiological analysis of soil samples from POME and non-POME sites. 2. To make recommendations on how best to handle POME in the study area. Methods: Quadrant approach was adopted for sampling POME (A) and Non POME (B) locations. Qualities were determined using standard analytical procedures. Conclusions: Results of the analysis were obtained in the following range; pH (3.940 -7.435), dissolved oxygen (DO) (1.582-6.234mg/l), biological oxygen demand (BOD) (50-5463mg/l etc. For the various locations, the population of total heterotrophic bacteria (THB) ranged from 1.36×10^6 - 2.42×10^6 cfu/ml, the total heterotrophic fungi (THF) ranged from 1.22 - 3.05×10^4 cfu/ml. The frequency of occurrence revealed the microbial isolates *Pseudomonas* sp., *Bacillus* sp., *Staphylococcus*, as the most frequently occurring isolates. Analysis of variance showed that there were significant differences ($P < 0.05$) in microbial populations among locations. The discharge of industrial effluents into the soil in Nigeria invariably results in the presence of high concentrations of pollutant in the soil environment.

Keywords : effluents, microbial composition, soil samples, isiala mbano

Conference Title : ICWMEE 2015 : International Conference on Waste Management and Environmental Engineering

Conference Location : London, United Kingdom

Conference Dates : February 16-17, 2015