

Does Mirror Therapy Improve Motor Recovery After Stroke? A Meta-Analysis of Randomized Controlled Trials

Authors : Hassan Abo Salem, Guo Feng, Xiaolin Huang

Abstract : The objective of this study is to determine the effectiveness of mirror therapy on motor recovery and functional abilities after stroke. The following databases were searched from inception to May 2014: Cochrane Stroke, Cochrane Central Register of Controlled Trials, MEDLINE, EMBASE, CINAHL, AMED, PsycINFO, and PEDro. Two reviewers independently screened and selected all randomized controlled trials that evaluate the effect of mirror therapy in stroke rehabilitation. 12 randomized controlled trials studies met the inclusion criteria; 10 studies utilized the effect of mirror therapy for the upper limb and 2 studies for the lower limb. Mirror therapy had a positive effect on motor recover and function; however, we found no consistent influence on activity of daily living, Spasticity and balance. This meta-analysis suggests that, Mirror therapy has additional effect on motor recovery but has a small positive effect on functional abilities after stroke. Further high-quality studies with greater statistical power are required in order to accurately determine the effectiveness of mirror therapy following stroke.

Keywords : mirror therapy, motor recovery, stroke, balance

Conference Title : ICNR 2015 : International Conference on Neurorehabilitation

Conference Location : Copenhagen, Denmark

Conference Dates : June 11-12, 2015