

Gastrointestinal Disturbances in Postural Orthostatic Tachycardia Syndrome (POTS)

Authors : Chandralekha Ashangari, Amer Suleman

Abstract : Background and Purpose: The Postural Orthostatic Tachycardia Syndrome (POTS) affects primarily young women. POTS is a form of dysautonomia that is estimated to impact between 1,000,000 and 3,000,000 Americans, and millions more around the world. POTS is a form of orthostatic intolerance that is associated with many Gastrointestinal disturbances. The aim of this study is to determine the Gastrointestinal disturbances in Postural Orthostatic Tachycardia Syndrome (POTS) patients.2. Methods: 249 patients referred to our clinic from January to November with POTS. Reviewed the medical records of 249 POTS patients and gastrointestinal symptoms. Results: however out of 249 patients, 226 patients are female (90.76%; average age 32.69), 23 patients are male (9.24%; average age 27.91) Data analysis: Out of 249 patients 189 patients (76%) had vomiting or nausea, 150 patients (60%) had irritable bowel syndrome, 128 patients (51%) had bloating, 125 patients (50%) had constipation , 80 patients (32%) had abdominal pain, 56 patients (22%) had delayed gastric emptying, 24 patients (10%) had lactose intolerance, 8 patients (3%) had Gastroesophageal reflux disease, 5 patients (2%) had Iron deficiency anemia, 6 patients (2%) had Peptic ulcer disease, 4 patients (2%) had Celiac Disease. Conclusion: Patients with POTS have a very high prevalence of gastrointestinal symptoms however the majority of abnormalities appear to be motility related. Motility testing should be performed in POTS patients. The diagnostic yield of endoscopic procedures appears to be low.

Keywords : gastrointestinal disturbances, Postural Orthostatic Tachycardia Syndrome (POTS), celiac disease, POTS patients

Conference Title : ICGU 2015 : International Conference on Gastroenterology and Urology

Conference Location : Miami, United States

Conference Dates : March 09-10, 2015