

## Utilization of Cervical Cancer Screening Among HIV Infected Women in Nairobi, Kenya

**Authors :** E. Njuguna, S. Ilovi, P. Muiruri, K. Mutai, J. Kinuthia, P. Njoroge

**Abstract :** Introduction: Cervical cancer is the commonest cause of cancer-related morbidity and mortality among women in developing countries in Sub Saharan Africa. Screening for cervical cancer in all women regardless of HIV status is crucial for the early detection of cancer of the cervix when treatment is most effective in curing the disease. It is particularly more important to screen HIV infected women as they are more at risk of developing the disease and progressing faster once infected with HPV (Human Papilloma Virus). We aimed to determine the factors affecting the utilization of cervical cancer screenings among HIV infected women above 18 years of age at Kenyatta National Hospital (KNH) Comprehensive Care Center (CCC). Materials and Methods: A cross-sectional mixed quantitative and qualitative study involving randomly and purposefully selected HIV positive female respectively was conducted. Qualitative data collection involved 4 focus group discussions of eligible female participants while quantitative data were acquired by one to one interviewer administered structured questionnaires. The outcome variable was the utilization of cervical cancer screening. Data were entered into Access data base and analyzed using Stata version 11.1. Qualitative data were analyzed after coding for significant clauses and transcribing to determine themes arising. Results: We enrolled a total of 387 patients, mean age (IQ range) 40 years (36-44). Cervical cancer screening utilization was 46% despite a health care provider recommendation of 85%. The screening results were reported as normal in 72 of 81 (88.9%) and abnormal 7 of 81(8.6%) of the cases. Those who did not know their result were 2 of 81(2.5%). Patients were less likely to utilize the service with increasing number of years attending the clinic (OR 0.9, 95% CI 0.86-0.99, p-value 0.02), but more likely to utilize the service if recommendation by a staff was made (OR 10, 95% CI 4.2-23.9, p<0.001), and if cervical screening had been done before joining KNH CCC (OR 2.9, 95% CI 1.7-4.9, p < 0.001). Similarly, they were more likely to rate the services on cervical cancer screening as good (OR 5.0, 95% CI 1.7-3.4, p <0.001) and very good (OR 8.1, 95% CI 2.5-6.1, p<0.001) if they had utilized the service. The main barrier themes emerging from qualitative data included fear of screening due to excessive pain or bleeding, lack of proper communication on screening procedures and increased waiting time. Conclusions: Utilization of cervical cancer screening services was low despite health care recommendation. Patient socio-demographic characteristics did not influence whether or not they utilized the services, indicating the important role of the health care provider in the referral and provision of the service.

**Keywords :** cervical, cancer, HIV, women, comprehensive care center

**Conference Title :** ICHA 2015 : International Conference on HIV and AIDS

**Conference Location :** London, United Kingdom

**Conference Dates :** May 25-26, 2015