

Biotransformation Process for the Enhanced Production of the Pharmaceutical Agents Sakuranetin and Genkwanin: Poised to be Potent Therapeutic Drugs

Authors : Niranjan Koirala, Sumangala Darsandhari, Hye Jin Jung, Jae Kyung Sohng

Abstract : Sakuranetin, an antifungal agent and genkwanin, an anti-inflammatory agent, are flavonoids with several potential pharmaceutical applications. To produce such valuable flavonoids in large quantity, an *Escherichia coli* cell factory has been created. *E. coli* harboring O-methyltransferase (SaOMT2) derived from *Streptomyces avermitilis* was employed for regiospecific methylation of naringenin and apigenin. In order to increase the production via biotransformation, metK gene was overexpressed and the conditions were optimized. The maximum yield of sakuranetin and genkwanin under optimized conditions was 197 μM and 170 μM respectively when 200 μM of naringenin and apigenin were supplemented in the separate cultures. Furthermore, sakuranetin was purified in large scale and used as a substrate for in vitro glycosylation by YjiC to produce glucose and galactose derivatives of sakuranetin for improved solubility. We also found that unlike naringenin, sakuranetin effectively inhibits α -melanocyte stimulating hormone (α -MSH)-stimulated melanogenesis in B16F10 melanoma cells. In addition, genkwanin more potently inhibited angiogenesis than apigenin. Based on our findings, we speculate that these compounds warrant further investigation in vivo as potential new therapeutic anti-carcinogenic, anti-melanogenic and anti-angiogenic agents.

Keywords : anti-carcinogenic, anti-melanogenic, glycosylation, methylation

Conference Title : ICBBB 2015 : International Conference on Bioscience, Biotechnology, and Biochemistry

Conference Location : Amsterdam, Netherlands

Conference Dates : August 06-07, 2015