

Determinants of Never Users of Contraception-Results from Pakistan Demographic and Health Survey 2012-13

Authors : Arsalan Jabbar, Wajiha Javed, Nelofer Mehboob, Zahid Memon

Abstract : Introduction: There are multiple social, individual and cultural factors that influence an individual's decision to adopt family planning methods especially among non-users in patriarchal societies like Pakistan. Non-users, if targeted efficiently, can contribute significantly to country's CPR. A research study showed that non-users if convinced to adopt lactational amenorrhea method can shift to long-term methods in future. Research shows that if non-users are targeted efficiently a 59% reduction in unintended pregnancies in Saharan Africa and South-Central and South-East Asia is anticipated. Methods: We did secondary data analysis on Pakistan Demographic Health Survey (2012-13) dataset. Use of contraception (never-use/ever-use) was the outcome variable. At univariate level Chi-square/Fisher Exact test was used to assess relationship of baseline covariates with contraception use. Then variables to be incorporated in the model were checked for multi-collinearity, confounding, and interaction. Then binary logistic regression (with an urban-rural stratification) was done to find the relationship between contraception use and baseline demographic and social variables. Results: The multivariate analyses of the study showed that younger women (≤ 29 years) were more prone to be never users as compared to those who were > 30 years and this trend was seen in urban areas (AOR 1.92, CI 1.453-2.536) as well as rural areas (AOR 1.809, CI 1.421-2.303). While looking at regional variation, women from urban Sindh (AOR 1.548, CI 1.142-2.099) and urban Balochistan (AOR 2.403, CI 1.504-3.839) had more never users as compared to other urban regions. Women in the rich wealth quintile were more never users and this was seen both in urban and rural localities (urban (AOR 1.106 CI .753-1.624); rural areas (AOR 1.162, CI .887-1.524)) even though these were not statistically significant. Women idealizing more children (> 4) are more never users as compared to those idealizing less children in both urban (AOR 1.854, CI 1.275-2.697) and rural areas (AOR 2.101, CI 1.514-2.916). Women who never lost a pregnancy were more inclined to be non-users in rural areas (AOR 1.394, CI 1.127-1.723). Women familiar with only traditional or no method had more never users in rural areas (AOR 1.717, CI 1.127-1.723) but in urban areas it wasn't significant. Women unaware of Lady Health Worker's presence in their area were more never users especially in rural areas (AOR 1.276, CI 1.014-1.607). Women who did not visit any care provider were more never users (urban (AOR 11.738, CI 9.112-15.121) rural areas (AOR 7.832, CI 6.243-9.826)). Discussion/Conclusion: This study concluded that government, policy makers and private sector family planning programs should focus on the untapped pool of never users (younger women from underserved provinces, in higher wealth quintiles, who desire more children.). We need to make sure to cover catchment areas where there are less LHWs and less providers as ignorance to modern methods and never been visited by an LHW are important determinants of never use. This all is in sync with previous literature from similar developing countries.

Keywords : contraception, demographic and health survey, family planning, never users

Conference Title : ICPD 2015 : International Conference on Population and Development

Conference Location : Montreal, Canada

Conference Dates : May 11-12, 2015