

Drug Sensitivity Pattern of Organisms Causing Chronic Suppurative Otitis Media

Authors : Fatma M. Benrabha

Abstract : The aim of the study was to determine the type and pattern of antibiotic susceptibility of the pathogenic microorganisms causing chronic suppurative otitis media (CSOM), which could lead to better therapeutic decisions and consequently avoidance of appearance of resistance to specific antibiotics. Most frequently isolated agents were *Pseudomonas aeruginosa* 28.5%; followed by *Staphylococcus aureus* 18.2%; *proteus mirabilis* 13.9%; *Providencia stuarti* 6.7%; *Bacteroides melaninogenicus*, *Aspergillus sp.*, *candida sp.*, 4.2% each; and other microorganisms were represented in 3-0.2%. Drug sensitivities pattern of *Pseudomonas aeruginosa* showed that ciprofloxacin was active against the majority of isolates (93.9%) followed by ceftazidime 86.2%, amikacin 76.2% and gentamicin 40.8%. However, *Staphylococcus aureus* isolates were resistant to penicillin 72.7%, erythromycin 28.6%, cephalothin 18.2%, cloxacillin 8.3% and ciprofloxacin was active against 96.2% of isolates. The resistance pattern of *proteus mirabilis* were 55.6% to ampicillin, 47.1% to carbencillin, 29.4% to cephalothin, 14.3% to gentamicin and 4.8% to amikacin while 100% were sensitive to ciprofloxacin. We conclude that ciprofloxacin is the best drug of choice in treatment of CSOM caused by the common microorganisms.

Keywords : otitis media, chronic suppurative otitis media (CSOM), microorganism, drug sensitivity

Conference Title : ICPP 2014 : International Conference on Pharmacy and Pharmacology

Conference Location : Bangkok, Thailand

Conference Dates : December 24-25, 2014