

Anthropometric Data Variation within Gari-Frying Population

Authors : T. M. Samuel, O. O. Aremu, I. O. Ismaila, L. I. Onu, B. O. Adetifa, S. E. Adegbite, O. O. Olokoshe

Abstract : The imperative of anthropometry in designing to fit cannot be overemphasized. Of essence is the variability of measurements among population for which data is collected. In this paper anthropometric data were collected for the design of gari-frying facility such that work system would be designed to fit the gari-frying population in the Southwestern states of Nigeria comprising Lagos, Ogun, Oyo, Osun, Ondo, and Ekiti. Twenty-seven body dimensions were measured among 120 gari-frying processors. Statistical analysis was performed using SPSS package to determine the mean, standard deviation, minimum value, maximum value and percentiles (2nd, 5th, 25th, 50th, 75th, 95th, and 98th) of the different anthropometric parameters. One sample t-test was conducted to determine the variation within the population. The 50th percentiles of some of the anthropometric parameters were compared with those from other populations in literature. The correlation between the worker's age and the body anthropometry was also investigated. The mean weight, height, shoulder height (sitting), eye height (standing) and eye height (sitting) are 63.37 kg, 1.57 m, 0.55 m, 1.45 m, and 0.67 m respectively. Result also shows a high correlation with other populations and a statistically significant difference in variability of data within the population in all the body dimensions measured. With a mean age of 42.36 years, results shows that age will be a wrong indicator for estimating the anthropometry for the population.

Keywords : anthropometry, cassava processing, design to fit, gari-frying, workstation design

Conference Title : ICHFE 2015 : International Conference on Human Factors and Ergonomics

Conference Location : Amsterdam, Netherlands

Conference Dates : August 06-07, 2015