## **Assessment of Dietary Intake of Pregnant Women**

Authors : Tuleshova Gulnara, Abduldayeva Aigul

**Abstract :** The goal is based on the studying the prevalence of micronutrient deficiencies among children and women of reproductive age to develop evidence-based recommendations aimed at improving the effectiveness of programs to prevent micronutrient deficiency. Subject: In our study we used a representative, random sample, carried out with the cluster method in the precinct of the principle areas of medical care for children 5 years of old. If the site has at least 60 children under 5 years of old, each second child was sampled, and if more than 60 children - each third child (first child selected by random sampling). The total number of investigated persons was within 80-86 women of reproductive age and children - within 80-92 people. Results: The studies found that the average prevalence of anemia among children aged 6-59 months was 35.2%, with the most susceptible to iron deficiency anemia in infants aged 6-23 months (53.3%). The prevalence of anemia among non-pregnant women was 39.0% among pregnant women - 43.8%. In children, the prevalence of folate deficiency was the highest (27.6%). Among non-pregnant women, frequent prevalence of folic acid deficiency was 37.0%. The prevalence of vitamin A deficiency was higher among children living in Astana (37.4%) compared with the medium-republican level (23.2%). **Keywords :** nutrition, pregnant women, micronutrients, macronutrients

Conference Title : ICEBB 2015 : International Conference on Environmental, Biomedical and Biotechnology

Conference Location : Venice, Italy

Conference Dates : November 09-10, 2015

1