

Sero-Prevalence of Hepatitis B Surface Antigen and Associated Factors among Pregnant Mothers Attending Antenatal Care Service, Mekelle, Ethiopia: Evidence from Institutional Based Quantitative Cross-Sectional Study

Authors : Semaw A., Awet H., Yohannes M.

Abstract : Background: Hepatitis B Virus (HBV) is a major global public health problem. Individuals living in Sub-Sahara Africa have 60% lifetime risk of acquiring HBV infection. Evidences showed that 80-90% of those born from infected mothers developed chronic HBV. Perinatal HBV transmission is a major determinant of HBV carrier status, its chronic sequel and maintains HBV transmission across generations. Method: Institution based cross-sectional study was conducted among 406 pregnant mothers attending Antenatal clinics at Mekelle and Ayder referral hospital from January 30 to April 1/2014. Epidata version 3.1 was used for data entry and SPSS version 21 statistical software was used for data cleaning, management and finally determine associated factors of hepatitis B surface antigen adjusting important confounders using multivariable logistic regression analysis at 5% level of significance. Result: The overall prevalence of hepatitis B surface antigen among pregnant women was 33 (8.1%). The socio-demographic characteristic of the study population showed that there is high positivity among secondary school 189 (46.6%). In the multivariable logistic regression analysis, history of a contact with individuals who had history of hepatitis B infection or jaundice and lifetime number of multiple sexual partners were found to be significantly associated with HBsAg positivity at AOR = 3.73 95%CI (1.373-10.182) and AOR = 2.57 95%CI (1.173-5.654), respectively. Moreover, Human Immunodeficiency Virus (HIV) and HBV confection rate was found 3.6%. Conclusion: This study has shown that HBV prevalence in pregnant women is highly prevalent (8.1%) in the study area. Contact with individuals who had a history of hepatitis or have jaundice and report of multiple lifetime sexual partnership were associated with hepatitis B infection. Education about HBV transmission and prevention as well as screening all pregnant mothers shall be sought to reduce the serious public health crisis of HBV.

Keywords : HBsAg, hepatitis B, pregnant women, prevalence

Conference Title : ICVH 2015 : International Conference on Viral Hepatitis

Conference Location : London, United Kingdom

Conference Dates : September 25-26, 2015