

## **Bone Marrow Edema Syndrome in the Foot and Ankle**

**Authors :** S. Alireza Mirghasemi, Elly Trepman, Mohammad Saleh Sadeghi, Narges Rahimi Gabaran, Shervin Rashidinia

**Abstract :** Bone marrow edema syndrome (BMES) is an uncommon and self-limited syndrome characterized by atraumatic extremity pain with unknown of etiology. Symptom onset may include sudden or gradual swelling and pain at rest or during activity, usually at night. This syndrome mostly affects middle-aged men and younger women who have pain in the lower extremities. The most common sites involved with BMES, in decreasing order of frequency, are the bones about the hip, knee, ankle, and foot. The diagnosis of BMES is made with magnetic resonance imaging to exclude other causes of bone marrow edema. The correct diagnosis often is delayed because of the low prevalence and nonspecific signs in the foot and ankle. This delay may intensify bone pain and impair patient function and quality of life. The goal of BMES treatment is to relieve pain and shorten disease duration. Treatment options are limited and may include symptomatic treatment, pharmacologic treatment, and surgery.

**Keywords :** transient osteoporosis, bone marrow edema syndrome, iloprost, bisphosphonates

**Conference Title :** ICOT 2015 : International Conference on Orthopedics and Traumatology

**Conference Location :** Venice, Italy

**Conference Dates :** August 13-14, 2015