Dimension of Water Accessibility in the Southern Part of Niger State, Nigeria

Authors : Kudu Dangana, Pai H. Halilu, Osesienemo R. Asiribo-Sallau, Garba Inuwa Kuta

Abstract : The study examined the determinants of household water accessibility in Southern part of Niger State, Nigeria. Data for the study was obtained from primary and secondary sources using questionnaire, interview, personal observation and documents. 1,192 questionnaires were administered; sampling techniques adopted are combination of purposive, stratified and simple random. Purposive sampling technique was used to determine sample frame; sample unit was determined using stratified sampling method and simple random technique was used in administering questionnaires. The result was analyzed within the scope of "WHO" water accessibility indicators using descriptive statistics. Major sources of water in the area are well; hand and electric pump borehole and streams. These sources account for over 90% of household's water. Average per capita water consumption in the area is 22 liters per day, while location efficiency of facilities revealed an average of 80 people per borehole. Household water accessibility is affected mainly by the factors of distances, time spent to obtain water, low income status of the majority of respondents to access modern water infrastructure, and to a lesser extent household size. Recommendations includes, all tiers of government to intensify efforts in providing water infrastructures and existing ones through budgetary provisions, and communities should organize fund raising bazaar, so as to raise fund to improve water infrastructures in the area.

1

Keywords : accessibility, determined, stratified, scope

Conference Title : ICEES 2016 : International Conference on Environmental Earth Sciences

Conference Location : Tokyo, Japan

Conference Dates : May 26-27, 2016