Comparative Antihyperglycemic Activity of Serpentina (Andrographis paniculata) and Papait (Mollugo oppositifolia linn) Aqueous Extracts in Alloxan-Induced Diabetic Mice

Authors : Karina Marie G. Nicolas, Kimberly M. Visaya, Emmanuel R. Cauinian

Abstract : A comparative study on the antihyperglycemic activity of aqueous extracts of Serpentina (Andrographis paniculata) and Papait (Mollugo oppositifolia linn) administered at 400mg/kg body weight per orem twice daily for 14 days was investigated using 24 alloxan-induced diabetic male, 6-8 weeks old ICR mice and Metformin as standard control. The blood glucose levels of all the animals in the treatment groups were not reduced to < 200mg/dl so as to consider them as non-diabetic but Papait showed a consistent blood glucose lowering effect from day 0 to 14 causing 36.07% reduction as compared to Serpentina which was observed to cause a fluctuating effect on blood glucose levels and a reduction of only 22.53% while the Metformin treated animals exhibited the highest reduction at 45.29%. The blood glucose levels at day 14 of animals treated with Papait (322.93 mg/dl) had comparable blood glucose levels (p<0.05) with the Metformin treated groups (284.50 mg/dl). Also, all the animals in the three treatment groups were still hypercholesterolemic with an observed consistent weight loss and a decrease in feed intake except for Serpentina which recorded a slight increase. Results of the study showed a superior antihyperglycemic activity of Papait compared with Serpentina.

Keywords : antihyperglycemic, diabetes, hypercholesterolemic, papait, serpentina

Conference Title : ICASVM 2015 : International Conference on Animal Science and Veterinary Medicine

Conference Location : Bangkok, Thailand

Conference Dates : December 17-18, 2015