

An Exploration Survival Risk Factors of Stroke Patients at a General Hospital in Northern Taiwan

Authors : Hui-Chi Huang, Su-Ju Yang, Ching-Wei Lin, Jui-Yao Tsai, Liang-Yiang

Abstract : Background: The most common serious complication following acute stroke is pneumonia. It has been associated with the increased morbidity, mortality, and medical cost after acute stroke in elderly patients. Purpose: The aim of this retrospective study was to investigate the relationship between stroke patients, risk factors of pneumonia, and one-year survival rates in a group of patients, in a tertiary referral center in Northern Taiwan. Methods: From January 2012 to December 2013, a total of 1730 consecutively administered stroke patients were recruited. The Survival analysis and multivariate regression analyses were used to examine the predictors for the one-year survival in stroke patients of a stroke registry database from northern Taiwan. Results: The risk of stroke mortality increased with age \geq 75 (OR=2.305, $p < .0001$), cancer (OR=3.221, $p < .0001$), stayed in intensive care unit (ICU) (OR=2.28, $p < .0006$), dysphagia (OR=5.026, $p < .0001$), without speech therapy (OR=0.192, $p < .0001$), serum albumin < 2.5 (OR=0.322, $p = .0053$), eGFR > 60 (OR=0.438, $p < .0001$), admission NIHSS > 11 (OR=1.631, $p = .0196$), length of hospitalization (d) > 30 (OR=0.608, $p = .0227$), and stroke subtype (OR=0.506, $p = .0032$). After adjustment of confounders, pneumonia was not significantly associated with the risk of mortality. However, it is most likely to develop in patients who are age ≥ 75 , dyslipidemia, coronary artery disease, albumin < 2.5 , eGFR < 60 , ventilator use, stay in ICU, dysphagia, without speech therapy, urinary tract infection, Atrial fibrillation, Admission NIHSS > 11 , length of hospitalization > 30 (d), stroke severity (mRS=3-5), stroke Conclusion: In this study, different from previous research findings, we found that elderly age, severe neurological deficit and rehabilitation therapy were significantly associated with Post-stroke Pneumonia. However, specific preventive strategies are needed to target the high risk groups to improve their long-term outcomes after acute stroke. These findings could open new avenues in the management of stroke patients.

Keywords : stroke, risk, pneumonia, survival

Conference Title : ICN 2016 : International Conference on Nursing

Conference Location : London, United Kingdom

Conference Dates : September 29-30, 2016