

## Prevalence of Autoimmune Thyroid Disease in Recurrent Aphthous Stomatitis

**Authors :** Arghavan Tonkaboni, Shamsolmolouk Najafi, Mohmmad Taghi Kiani, Mehrzad Gholampour, Touraj Goli

**Abstract :** Introduction: Recurrent aphthous stomatitis (RAS) is a multifactorial recurrent oral lesion; which is an autoimmune disease. TH1 cytokines are the most important etiological factors. Autoimmune thyroid disease (ATD) is one of the most common autoimmune diseases and generally coexists with other autoimmune diseases. This study assessed the prevalence of thyroid disease in patients with recurrent aphthous stomatitis. Materials and Methods: This case control study assessed 100 known RAS patients who were diagnosed clinically by oral medicine specialists; venous blood samples were analyzed for thyroid stimulating hormone (TSH), free triiodothyronine (fT3), total thyroxine (fT4), thyroglobulin, anti-thyroid peroxidase antibody (anti-TPO) and anti-thyroglobulin antibody (anti-TG) levels. Results: Fifty patients with RAS aged between 18-42 years ( $28.5 \pm 5.8$ ) and 50 healthy volunteers aged 19-45 years ( $27.3 \pm 5.4$ ) participated. In RAS patients, fT3 and TSH levels were significantly higher ( $P=0.031$ ,  $P=0.706$ ); however, fT4 level was lower in the RAS group ( $P=0.447$ ). Anti TG and anti-TPO levels were significantly higher in the RAS group ( $P=0.008$ ,  $P=0.067$ ). Conclusion: Our study showed that ATD prevalence was significantly higher in RAS patients. Based on this study, we recommend assessment of thyroid hormones and antibodies in RAS patients.

**Keywords :** recurrent aphthous stomatitis, thyroid antibodies, thyroid hormone, thyroid autoimmune disease

**Conference Title :** ICVID 2016 : International Conference on Virology and Infectious Diseases

**Conference Location :** London, United Kingdom

**Conference Dates :** February 25-26, 2016