

Association Nephropathy and Hypertension in Diabetic Patients

Authors : Bahlous Afef, Bouzid Kahena, Bardkis Ahlem, Mrad Mehdi, Kalai Eya, Sonia Bahri, Abdelmoula Jaouida

Abstract : Diabetic nephropathy is the first cause of chronic renal failure and hemodialysis use in several countries including Tunisia. The role of hypertension (HT) as major risk factor for nephropathy is undeniable. The aim of our study was to determine the relationship between blood pressure and nephropathy in a population of diabetic type 2 recently discovered. Materials and methods: We conducted a prospective study focused on 60 patients with type 2 diabetes recently discovered (<5 years). Each patient have benefited from: -a full clinical examination with measurement of blood pressure - exploring a blood-glucose control and renal function -urinary exploration with the determination of proteinuria microalbuminurie of 24 hours with a immunoturbidimetric method using Architect (ABBOTT CI 8200). Results and discussion: Hypertension was present in 46.7% of cases. Twenty patients, 35% of the study population showed nephropathy. Four of these patients (6.66% of cases) had proteinuria, while 16 (26.6% of patients) had microalbuminuria (> 30mg/24 hours). Systolic blood pressure was significantly ($p < 0.05$) associated with the presence of nephropathy (139 ± 19.44) vs. for the group with normal renal function (128.65 ± 15.12 mmHg). Conclusion: The etiology of diabetic nephropathy is multifactorial. However, systolic blood pressure and glyceemic control remains the major risk factors. Better glyceemic control and treatment of hypertension allowed preventing and slowing the progression of diabetic nephropathy.

Keywords : hypertension, nephropathy, hemodialysis, diabetes

Conference Title : ICPHCMS 2016 : International Conference on Prehypertension, Hypertension and Cardio Metabolic Syndrome

Conference Location : Venice, Italy

Conference Dates : April 11-12, 2016