

Detection of Heroin and Its Metabolites in Urine Samples: A Chemiluminescence Approach

Authors : Sonu Gandhi, Neena Capalash, Prince Sharma, C. Raman Suri

Abstract : A sensitive chemiluminescence immunoassay (CIA) for heroin and its major metabolites is reported. The method is based on the competitive reaction of horseradish peroxidase (HRP)-labeled anti-MAM antibody and free drug in spiked urine samples. A hapten-protein conjugate was synthesized by using acidic derivative of monoacetyl morphine (MAM) coupled to carrier protein BSA and was used as an immunogen for the generation of anti-MAM (monoacetyl morphine) antibody. A high titer of antibody (1:64,000) was obtained and the relative affinity constant (K_{aff}) of antibody was 3.1×10^7 l/mol. Under the optimal conditions, linear range and reactivity for heroin, mono acetyl morphine (MAM), morphine and codeine were 0.08, 0.09, 0.095 and 0.092 ng/mL respectively. The developed chemiluminescence inhibition assay could detect heroin and its metabolites in standard and urine samples up to 0.01 ng/ml.

Keywords : heroin, metabolites, chemiluminescence immunoassay, horse radish peroxidase

Conference Title : ICB 2016 : International Conference on Biosensors

Conference Location : San Francisco, United States

Conference Dates : June 09-10, 2016