

## Antibacterial and Antityrosinase Activity of Isolated Compounds from Stem Bark of *Ficus platyphylla* Del

**Authors :** Aminu Muhammad, Mustapha Ya'u, Hasnah Mohd Sirat

**Abstract :** An investigation of the chemical constituents into the stem bark of *Ficus platyphylla* (Moraceae) has resulted in the isolation of hordenine, epicatechin, lupeol, lupeol acetate and  $\alpha$ -amyirin acetate. Their structures were determined using spectroscopic data as well as comparison with literature data. The antibacterial assay has been tested against Gram positive and Gram negative bacteria, while the tyrosinase inhibition assay was examined using L-Dopa as a substrate of mushroom tyrosinase enzyme. hordenine, epicatechin, lupeol, lupeol acetate and  $\alpha$ -amyirin acetate showed minimum inhibition concentration (MIC) values in the range of 225-900  $\mu\text{g/mL}$  against the bacterial strains. Lupeol, lupeol acetate and  $\alpha$ -amyirin acetate showed significant antityrosinase activity against mushroom tyrosinase enzyme with percent inhibition of 67.7%, 66.2% and 62.2%, respectively.

**Keywords :** antibacterial, antityrosinase, chemical constituents, *Ficus platyphylla*

**Conference Title :** ICSRD 2020 : International Conference on Scientific Research and Development

**Conference Location :** Chicago, United States

**Conference Dates :** December 12-13, 2020