

Length-Weight and Length-Length Relationships for 14 Sparidae Species, from the Northeastern Mediterranean Sea Coast of Turkey

Authors : Hacer Yeldan, Erhan Akamca, Sedat Gündoğdu

Abstract : Length-Weight and Length-length relationship were estimated of 14 species Sparidae (Boops boops, Diplodus annularis, Diplodus cervinus, Diplodus puntazzo, Diplodus sargus, Diplodus vulgaris, Lithognathus mormyrus, Oblada melanura, Pagellus acarne, Pagellus erythrinus, Pagrus auriga, Pagrus caeruleostictus, Sarpa salpa, Sparus aurata) sampled from in the Northeastern Mediterranean Sea coast of Turkey, Iskenderun Bay. Samples were collected from July 2014 to June 2015, using bottom trawl and trammel net into three different depth; 0-10 m, 10-20 m, 20-50m. Length-length relationships were determined size measurements: standard length (SL) and fork length (FL) to total length (TL) for fish species. The relationships between TL, FL and TL, SL were all linear. The values of the exponent b of the length-weight relationships ranged between 2.685 and 3.473. The type of growth for fish species was allometric growth.

Keywords : Sparidae, Iskenderun bay, length-length, length-weight relationships

Conference Title : ICEAFS 2016 : International Conference on Environment, Agriculture and Food Sciences

Conference Location : Lisbon, Portugal

Conference Dates : April 14-15, 2016