

## Progressive Type-I Interval Censoring with Binomial Removal-Estimation and Its Properties

**Authors :** Sonal Budhiraja, Biswabrata Pradhan

**Abstract :** This work considers statistical inference based on progressive Type-I interval censored data with random removal. The scheme of progressive Type-I interval censoring with random removal can be described as follows. Suppose  $n$  identical items are placed on a test at time  $T_0 = 0$  under  $k$  pre-fixed inspection times at pre-specified times  $T_1 < T_2 < \dots < T_k$ , where  $T_k$  is the scheduled termination time of the experiment. At inspection time  $T_i$ ,  $R_i$  of the remaining surviving units  $S_i$ , are randomly removed from the experiment. The removal follows a binomial distribution with parameters  $S_i$  and  $p_i$  for  $i = 1, \dots, k$ , with  $p_k = 1$ . In this censoring scheme, the number of failures in different inspection intervals and the number of randomly removed items at pre-specified inspection times are observed. Asymptotic properties of the maximum likelihood estimators (MLEs) are established under some regularity conditions. A  $\beta$ -content  $\gamma$ -level tolerance interval (TI) is determined for two parameters Weibull lifetime model using the asymptotic properties of MLEs. The minimum sample size required to achieve the desired  $\beta$ -content  $\gamma$ -level TI is determined. The performance of the MLEs and TI is studied via simulation.

**Keywords :** asymptotic normality, consistency, regularity conditions, simulation study, tolerance interval

**Conference Title :** ICRM 2016 : International Conference on Reliability and Maintainability

**Conference Location :** Barcelona, Spain

**Conference Dates :** August 11-12, 2016