

## **A Paradox in the Issue of Sexual Violence: A Study on Sexual Violence Perpetrated against Men and Boys by Women: A Case Study of the Municipality of Ibanda, Town of Bukavu, Province of South Kivu, Democratic Republic of Congo, Africa**

**Authors :** Sylvie Ekanga Lumumba

**Abstract :** Background and Significance of the Study: Over the past three decades, the perception of sexual violence has changed significantly, it is now recognized that men and boys are victims of sexual violence. However, the body of research on male victims and particularly on their attackers is much more limited. Research on the above is thus more than required. To contribute to the above quest for further studies, the researcher conducted this study on sexual violence perpetrated against men and boys by women, in the Municipality of Ibanda, Town of Bukavu, Province of South Kivu, Democratic Republic of Congo. The main study objectives were the following: to investigate on the statement of sexual violence perpetrated against men and boys in the Municipality of Ibanda, to investigate into its consequences and the statement of medical and psycho-social care given to victims. Methodology: Data were collected using valid and reliable Survey Questionnaire and Interview Schedule. Study population: the 85,882 men and boys from the Municipality of Ibanda. Sampling: led to 150 men and boys, received discreetly by the researcher during November-December 2015. Major findings: First, findings related to sexual abuse and its procedure: 74.2% of men and boys were victims of sexual violence perpetrated by a woman, more than a year ago. 13.3% however, were victims for less than a year now. 79.7% of victims have experienced sexual violence by a sexual act; 3.9% through the intention of the woman to cause the death of the victim, by serious injury to the genitals. The Second group of findings related to the consequences of sexual violence revealed that HIV/AIDS is the most important physical consequence experienced by 77.3 % of victims. Physical psychological consequences are: urinary or defecation problems (72.7%); while key psycho-emotional and behavioral consequence is: living a state of deep shame and humiliation: 68.8%. As for sexual consequences: 71.1% indicated a chronic avoidance of sexual activity and 57% reported sexual dysfunctions. The third group of findings is related to medical and psycho-social care: repetitively, more than 80% of male victims affirmed that with the help of friends and traditional healers, they took care of themselves for all the eight WHO phases of clinical care of rape victims, this was hence not effectively done. Concluding Statement: for this study, the statement of sexual violence of men and boys by women in the Eastern Congo and its consequences are not researched upon and are underestimated; the study also revealed that the care of male victims is grossly ill-conducted, as opposed to female victims care. It therefore calls for further research and further vulgarization of the research results, to convince other stakeholders (politicians for example) to immediately take action.

**Keywords :** sexual violence, men and boys, medical care, psycho-social care

**Conference Title :** ICGSDS 2016 : International Conference on Gender, Sexuality and Diversity Studies

**Conference Location :** Vancouver, Canada

**Conference Dates :** August 04-05, 2016