

Prevalence of *Cryptosporidium* spp. in Free-Living Wild Birds by Using Carbol Fuchsin Staining Methods in Konya, Turkey

Authors : Nermin Isik

Abstract : Cryptosporidiosis is one of the most common parasitic infection in domesticated, caged, wild birds. *Cryptosporidium* sp. has been reported in over 30 avian species worldwide. *Cryptosporidium meleagridis*, *Cryptosporidium baileyi* and *Cryptosporidium galli* are recognised avian species of *Cryptosporidium*. This study was carried out to determine the prevalence of *Cryptosporidium* sp. in wild birds in Konya province, Turkey. Faecal samples were collected from 65 wild birds including 52 Podicipedidae (*Podiceps cristatus*), 11 Rallidae (*Fulicia Atra*), 2 Anitadia (*Aythya ferina*). Faecal samples were stained with Modified Ziehl-Neelsen staining technique, they were examined under light microscope for the presence of *Cryptosporidium* sp. oocysts. Among the 65 faecal samples, 11 (16.9%) were found to be infected with *Cryptosporidium* sp. oocysts. The results of this study indicate that wild birds may play an important role in the epidemiology of *Cryptosporidium*. In conclusion, Cryptosporidiosis has suggested zoonotic potential and thus warrant further attention. In addition, biological and genetic studies are required to provide more information on Cryptosporidiosis.

Keywords : *Cryptosporidium* sp, wild birds, Konya, Turkey

Conference Title : ICAFBBE 2016 : International Conference on Agricultural Sciences, Food, Beverage and Bioprocess Engineering

Conference Location : Amsterdam, Netherlands

Conference Dates : May 12-13, 2016