

Effect of Nitrogen and Carbon Sources on Growth and Lipid Production from Mixotrophic Growth of *Chlorella* sp. KKU-S2

Authors : Ratanaporn Leasing, Thidarat Papone, Mutiyaporn Puangbut

Abstract : Mixotrophic cultivation of the isolated freshwater microalgae *Chlorella* sp. KKU-S2 in batch shake flask for biomass and lipid productions, different concentration of glucose as carbon substrate, different nitrogen source and concentrations were investigated. Using 1.0g/L of NaNO₃ as nitrogen source, the maximum biomass yield of 10.04g/L with biomass productivity of 1.673g/L d was obtained using 40g/L glucose, while a biomass of 7.09, 8.55 and 9.45g/L with biomass productivity of 1.182, 1.425 and 1.575g/L d were found at 20, 30 and 50g/L glucose, respectively. The maximum lipid yield of 3.99g/L with lipid productivity of 0.665g/L d was obtained when 40g/L glucose was used. Lipid yield of 1.50, 3.34 and 3.66g/L with lipid productivity of 0.250, 0.557 and 0.610g/L d were found when using the initial concentration of glucose at 20, 30 and 50g/L, respectively. Process product yield (YP/S) of 0.078, 0.119, 0.158 and 0.094 were observed when glucose concentration was 20, 30, 40 and 50 g/L, respectively. The results obtained from the study shows that mixotrophic culture of *Chlorella* sp. KKU-S2 is a desirable cultivation process for microbial lipid and biomass production.

Keywords : mixotrophic cultivation, microalgal lipid, *Chlorella* sp. KKU-S2

Conference Title : ICSRD 2020 : International Conference on Scientific Research and Development

Conference Location : Chicago, United States

Conference Dates : December 12-13, 2020