

Ethanol and Biomass Production from Spent Sulfite Liquor by Filamentous Fungi

Authors : M. T. Asadollahzadeh, A. Ghasemian, A. R. Saraeian, H. Resalati, P. R. Lennartsson, M. J. Taherzadeh

Abstract : Since filamentous fungi are capable of assimilating several types of sugars (hexoses and pentoses), they are potential candidates for bioconversion of spent sulfite liquor (SSL). Three filamentous fungi such as *Aspergillus oryzae*, *Mucor indicus*, and *Rhizopus oryzae* were investigated in this work. The SSL was diluted in order to obtain concentrations of 50, 60, 70, 80, and 90% and supplemented with two types of nutrients. The results from cultivations in shake flask showed that *A. oryzae* and *M. indicus* were not able to grow in pure SSL and SSL90% while *R. oryzae* could grow only in SSL50% and SSL60%. Cultivation with *A. oryzae* resulted in the highest yield of produced fungal biomass, while *R. oryzae* cultivation resulted in the lowest fungal biomass yield. Although, the mediums containing yeast extract, $(\text{NH}_4)_2\text{SO}_4$, KH_2PO_4 , $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, and $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ as nutrients supplementations produced higher fungal biomass compared to the mediums containing $\text{NH}_4\text{H}_2\text{PO}_4$ and ammonia, but there was no significant difference between two types of nutrients in terms of sugars and acetic acid consumption rate. The sugars consumption in *M. indicus* cultivation was faster than *A. oryzae* and *R. oryzae* cultivation. Acetic acid present in SSL was completely consumed during cultivation of all fungi. *M. indicus* was the best and fastest ethanol producer from SSL among the fungi examined, when yeast extract and salts were used as nutrients supplementations. Furthermore, no further improvement in ethanol concentration and rate of sugars consumption was obtained in medium supplemented with $\text{NH}_4\text{H}_2\text{PO}_4$ and ammonia compared to medium containing yeast extract, $(\text{NH}_4)_2\text{SO}_4$, KH_2PO_4 , $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, and $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$. On the other hand, the higher dilution of SSL resulted in a better fermentability, and better consumption of sugars and acetic acid.

Keywords : ethanol, filamentous fungi, fungal biomass, spent sulfite liquor

Conference Title : ICBB 2017 : International Conference on Biotechnology and Bioengineering

Conference Location : Paris, France

Conference Dates : October 19-20, 2017