## Effect of Varying Levels of Concentrate Ration on the Performance of Nili-Ravi Buffalo Heifer Calves

Authors : Z. M. Iqbal, M. Abdullah, K. Javed, M. A. Jabbar, A. Haque, M. Saadullah, F. Shahzad

**Abstract :** The current study was conducted to set the appropriate concentrate level for Nili-Ravi buffalo heifers. Twenty seven buffalo heifers were randomly divided into three different groups A, B and C having nine animals in each group. All the heifers were given free access to chopped green fodder and fresh water. In addition, heifers of group A, B and C were given concentrate at the rate of 0.5%, 1% and 1.5% of their body weight. The average daily dry matter intake was 2.69, 3.06 and 3.83 kg with average daily gain of 456.09, 398.56 and 515.87 gm in group A, B and C, respectively. The feed conversion ratio of heifers of these groups was 5.89, 7.74 and 7.52, respectively. There was non-significant (P>0.05) difference in the body measurements (height at wither, body length and heart girth), final body condition and scoring and blood serum (glucose, total protein and cholesterol) of heifers of all the three groups. The results of current study shows that there is non-significant (P>0.05) difference in the growth rate of Nili-Ravi heifers at varying levels of concentrate so, it is cost effective to raise 6-8 month calves by offering concentrate at the rate of 0.5% body weight along with free access of green fodder.

Keywords : concentrate level, buffalo heifer, body measurement, green fodder

Conference Title : ICAFAS 2014 : International Conference on Agricultural, Food and Animal Sciences

Conference Location : Zurich, Switzerland

Conference Dates : July 30-31, 2014