

## **Determination of Some Biochemical Parameters in Women during the First Trimester of Pregnancy (Normal Pregnancy and Missed Miscarriage)**

**Authors :** M. Yahia, N. Chaoui, A. Chaouch, Massinissa Yahia

**Abstract :** Our study was designed to determine the metabolic changes of some biochemical parameters (cholesterol, triglyceride, Iron, uric acid, Urea and folic acid) and highlight their changes in 57 women of the region Batna, during the first trimester of pregnancy. This practical work was done with 27 women with missed miscarriage, compared with 30 control subjects of normal pregnant women. The assay results revealed a highly significant difference ( $P = 0.0006$ ) between the two groups in serum iron (64.00 vs 93.54) and in the rate of folate (6.70 vs 9.22) ( $P < 0.001$ ) but no difference was found regarding the rate of Ca (9.69 vs 10.20), urea (0.19 vs 0.17), UA (33.96 vs 32.76), CH (1.283 vs 1.431), and TG (0.8852 vs 0.8290). The present study indicates that iron deficiency and folate are associated with missed miscarriage, but no direct pathophysiological link has been determined. Further in-depth studies are needed to determine the exact mechanism by which these deficits lead to a missed miscarriage.

**Keywords :** biochemical parameters, pregnant women, missed miscarriage, Algeria

**Conference Title :** ICABBBE 2014 : International Conference on Agricultural, Biotechnology, Biological and Biosystems Engineering

**Conference Location :** Barcelona, Spain

**Conference Dates :** February 27-28, 2014