

## Some Remains of Fossil Artiodactyla: Evolutionary Status, Taxonomy and Biogeographical Distribution in Late Miocene of Pakistan

**Authors :** Khizar Samiullah Samiullah, Riffat Yasin, Khurram Feroz, Omer Draz, Memmona Nazish

**Abstract :** New fossil remains of artiodactyl have been recovered from three Late Miocene localities, Lava, Dhok Bun Ameer Khatoon and Hasnoot. These localities belong to lower and middle Siwalik Hills of Pakistan, the Chinji and Dhok Pathan Formation respectively and are remarkably rich in fossils of artiodactyl. The fauna mainly comprises various families of order Artiodactyla; Cervidae, Equidae, Proboscidea, Giraffidae, Rhinocerotidae, Tragulidae, Suidae and Primates. In Chinji Formation Lava and Dhok Bun Ameer Khatoon are located in district Chakwal while in Upper Dhok Pathan Formation the best fossils exposure site is Hasnoot which is located in District Jhelum, Punjab, Pakistan. Specimens described and discussed here include right and left maxilla, isolated upper premolars and molars which have been collected during extensive fieldwork. After morphological and comparative analysis the collection is attributed to Giraffokeryx, Giraffa, Listriodon, Dorcatherium, Selenoportax and Pachyportax. In this study evolutionary status, taxonomy and biogeographical distribution as well as the relationship of different Artiodactyls have been discussed comprehensively. The Palaeoenvironmental studies reveal the persistence of mosaics of diverse habitats ranging from tropical evergreen forest to subtropical ones, closed seasonal woodlands to wooded savannas during the deposition of these outcrops.

**Keywords :** Artiodactyla, fossil dentition, late Miocene, lower and middle Siwaliks

**Conference Title :** ICSRD 2020 : International Conference on Scientific Research and Development

**Conference Location :** Chicago, United States

**Conference Dates :** December 12-13, 2020