

## Expression of Fused Plasmodium falciparum Orotate Phosphoribosyltransferase and Orotidine 5'-Monophosphate Decarboxylase in Escherichia coli

**Authors :** Waranya Imprasittichai, Patsarawadee Paojinda, Sudaratana R. Krungkrai, Nirianne Marie Q. Palacpac, Toshihiro Horii, Jerapan Krungkrai

**Abstract :** Fusion of the last two enzymes in the pyrimidine biosynthetic pathway in the inversed order by having COOH-terminal orotate phosphoribosyltransferase (OPRT) and NH<sub>2</sub>-terminal orotidine 5'-monophosphate decarboxylase (OMPDC), as OMPDC-OPRT, are described in many organisms. In this study, we constructed gene fusions of Plasmodium falciparum OMPDC-OPRT (1,836 bp) in pTrcHisA vector and expressed as an 6xHis-tag bifunctional protein in three Escherichia coli strains (BL21, Rosetta, TOP10) at 18 °C, 25 °C and 37 °C. The recombinant bifunctional protein was partially purified by Ni-Nitrilotriacetic acid-affinity chromatography. Specific activities of OPRT and OMPDC domains in the bifunctional enzyme expressed in E. coli TOP10 cells were approximately 3-4-fold higher than those in BL21 cells. There were no enzymatic activities when the construct vector expressed in Rosetta cells. Maximal expression of the fused gene was observed at 18 °C and the bifunctional enzyme had specific activities of OPRT and OMPDC domains in a ratio of 1:2. These results provide greater yields and better catalytic activities of the bifunctional OMPDC-OPRT enzyme for further purification and kinetic study.

**Keywords :** bifunctional enzyme, orotate phosphoribosyltransferase, orotidine 5'-monophosphate decarboxylase, plasmodium falciparum

**Conference Title :** ICABMB 2017 : International Conference on Advances in Biochemistry and Molecular Biology

**Conference Location :** Tokyo, Japan

**Conference Dates :** May 28-29, 2017