

Relationship between Body Mass Composition and Primary Dysmenorrhoea

Authors : Snehalata Tembhurne

Abstract : Introduction: A healthy menstrual cycle is a sign of women's sound health. Various variables may influence the length and regularity of menstrual cycle. Studies have revealed that menstrual cycle abnormalities may be associated with psychological stress, lack of physical exercise, alteration in body composition, endocrine disturbances, higher estrogen levels as seen in obese females. Hence there is an urgent need to find out the relationship between variations in body mass composition (BMI & body fat%) with menstrual abnormalities like primary dysmenorrhoea. Aim: To find out the relationship between body mass composition and primary dysmenorrhoea. Objectives: 1. To check whether there is any association between body mass index and primary dysmenorrhoea. 2. To check whether there is any association between body fat percentage and primary dysmenorrhoea. NULL HYPOTHESES- There is no relationship between body mass composition and primary dysmenorrhoea. Hypothesis: There exists a relationship between body mass composition and primary dysmenorrhoea. Materials and Methods: The study was conducted over a period of 6 months with 90 samples selected on random basis. The procedure was explained to the participant and a written consent was taken thereafter. The participant was made to stand on the BODY COMPOSITION SCANNING MONITOR, which scanned the physical profile of the participant (height, weight, BMI, body fat percentage and visceral fat). Thereafter, the candidate was asked about her menstrual irregularities and was asked to grade her level of dysmenorrhoea (if present) using the Verbal Dimensional Dysmenorrhoea Scale. Results: Chi square test of association was used to find out the association between body mass composition (body mass index, body fat percentage) and primary dysmenorrhoea. The chi-square value for association between body mass index and primary dysmenorrhoea was 38.63 $p < 0.001$ which was statistically significant. The chi-square value for the association of body fat % & primary dysmenorrhoea was 30.09, $p < 0.001$ which was statistically significant. Conclusion: Study shows that there exists a significant relationship between body mass composition and primary dysmenorrhoea and as the value of Body mass index and body fat percentages goes on increasing in females, the severity of primary dysmenorrhoea also increases.

Keywords : body mass index, body composition screening monitor, primary dysmenorrhoea, verbal dimensional dysmenorrhoea scale

Conference Title : ICSRD 2020 : International Conference on Scientific Research and Development

Conference Location : Chicago, United States

Conference Dates : December 12-13, 2020