

## **An Analytical Wall Function for 2-D Shock Wave/Turbulent Boundary Layer Interactions**

**Authors :** X. Wang, T. J. Craft, H. Iacovides

**Abstract :** When handling the near-wall regions of turbulent flows, it is necessary to account for the viscous effects which are important over the thin near-wall layers. Low-Reynolds- number turbulence models do this by including explicit viscous and also damping terms which become active in the near-wall regions, and using very fine near-wall grids to properly resolve the steep gradients present. In order to overcome the cost associated with the low-Re turbulence models, a more advanced wall function approach has been implemented within OpenFoam and tested together with a standard log-law based wall function in the prediction of flows which involve 2-D shock wave/turbulent boundary layer interactions (SWTBLIs). On the whole, from the calculation of the impinging shock interaction, the three turbulence modelling strategies, the Launder-Sharma k- $\epsilon$  model with Yap correction (LS), the high-Re k- $\epsilon$  model with standard wall function (SWF) and analytical wall function (AWF), display good predictions of wall-pressure. However, the SWF approach tends to underestimate the tendency of the flow to separate as a result of the SWTBLI. The analytical wall function, on the other hand, is able to reproduce the shock-induced flow separation and returns predictions similar to those of the low-Re model, using a much coarser mesh.

**Keywords :** SWTBLIs, skin-friction, turbulence modeling, wall function

**Conference Title :** ICCFWT 2017 : International Conference on Compressible Flows and Wave Turbulence

**Conference Location :** London, United Kingdom

**Conference Dates :** January 19-20, 2017