

Efficacy and Safety by Baseline A1c with Once-Weekly Dulaglutide in the AWARD Program

Authors : Alaa Mostafa, Samuel Dagogo-Jack, Vivian Thieu, Maria Yu, Nan Zhang, Dara Schuster, Luis-Emilio Garcia-Perez

Abstract : Dulaglutide (DU), a once-weekly glucagon-like peptide-1 receptor agonist, was studied in the AWARD clinical trial program in adult patients with type 2 diabetes (T2D) and demonstrated significant hemoglobin A1c (A1c) reduction and potential for weight loss. To evaluate the efficacy and safety of DU 1.5 mg and DU 0.75 mg in patients with T2D by baseline A1c <8.5% or ≥8.5%, a post-hoc analysis was conducted on AWARD-1 to -6 and -8 at 6 months. Across 7 studies, 55% to 82% of the DU-treated patients had a baseline A1c <8.5%, and 18% to 45% had a baseline A1c ≥8.5%. The ranges of A1c reductions with baseline A1c <8.5% and ≥8.5%, respectively, were: DU 1.5 mg: -0.67% to -1.25% and -1.22% to -2.37%; DU 0.75 mg: -0.53% to -1.07% and -1.37% to -2.19%. The A1c reduction from the pooled analysis was greater in patients with baseline A1c ≥8.5% than patients with baseline A1c <8.5%, respectively: DU 1.5 mg: -1.86% and -1.02%; DU 0.75 mg: -1.75% and -0.83%. DU treatments were well tolerated among baseline A1c subgroups. Across the AWARD program, DU 1.5 mg and DU 0.75 mg demonstrated significant A1c reduction in both subgroups with an acceptable safety profile. Compared to patients with baseline A1c <8.5%, patients with baseline A1c ≥8.5% had greater A1c reduction. Disclosures: This study was supported and conducted by Eli Lilly and Company, Indianapolis, IN, USA.

Keywords : A1c reduction, dulaglutide, type 2 diabetes, weight loss

Conference Title : ICCED 2017 : International Conference on Clinical Endocrinology and Diabetes

Conference Location : Jeddah, Saudi Arabia

Conference Dates : January 30-31, 2017