

## The Impact of Physical Activity for Recovering Cancer Patients

**Authors :** Martyn Queen, Diane Crone, Andrew Parker, Saul Bloxham

**Abstract :** Rationale: There is a growing body of evidence that supports the use of physical activity during and after cancer treatment. However, activity levels for patients remain low. As more cancer patients are treated successfully, and treatment costs continue to escalate, physical activity may be a promising adjunct to a person-centred healthcare approach to recovery. Aim: The aim was to further understand how physical activity may enhance the recovery process for a group of mixed-site cancer patients. Objectives: The research investigated longitudinal changes in physical activity and perceived the quality of life between two and six month's post-exercise interventions. It also investigated support systems that enabled patients to sustain these perceived changes. Method: The respondent cohort comprised 14 mixed-site cancer patients aged 43-70 (11 women, 3 men), who participated in a two-phase physical activity intervention that took place at a university in the South West of England. Phase 1 consisted of an eight-week structured physical activity programme; Phase 2 consisted of four months of non-supervised physical activity. Semi-structured interviews took place three times over six months with each participant. Grounded theory informed the data collection and analysis which, in turn, facilitated theoretical development. Findings: Our findings propose three theories on the impact of physical activity for recovering cancer patients: 1) Knowledge gained through a structured exercise programme can enable recovering cancer patients to independently sustain physical activity to four-month follow-up. 2) Sustaining physical activity for six months promotes positive changes in the quality of life indicators of chronic fatigue, self-efficacy, the ability to self-manage and energy levels. 3) Peer support from patients facilitates adherence to a structured exercise programme and support from a spouse, or life partner facilitates independently sustained physical activity to four-month follow-up. Conclusions: This study demonstrates that qualitative research can provide an evidence base that could be used to support future care plans for cancer patients. Findings also demonstrate that a physical activity intervention can be effective at helping cancer patients recover from the side effects of their treatment, and recommends that physical activity should become an adjunct therapy alongside traditional cancer treatments.

**Keywords :** physical activity, health, cancer recovery, quality of life, support systems, qualitative, grounded theory, person-centred healthcare

**Conference Title :** ICPAH 2017 : International Conference on Physical Activity and Health

**Conference Location :** Kuala Lumpur, Malaysia

**Conference Dates :** August 24-25, 2017