

Anxiety Treatment: Comparing Outcomes by Different Types of Providers

Authors : Melissa K. Hord, Stephen P. Whiteside

Abstract : With lifetime prevalence rates ranging from 6% to 15%, anxiety disorders are among the most common childhood mental health diagnoses. Anxiety disorders diagnosed in childhood generally show an unremitting course, lead to additional psychopathology and interfere with social, emotional, and academic development. Effective evidence-based treatments include cognitive-behavioral therapy (CBT) and selective serotonin reuptake inhibitors (SSRI's). However, if anxious children receive any treatment, it is usually through primary care, typically consists of medication, and very rarely includes evidence-based psychotherapy. Despite the high prevalence of anxiety disorders, there have only been two independent research labs that have investigated long-term results for CBT treatment for all childhood anxiety disorders and two for specific anxiety disorders. Generally, the studies indicate that the majority of youth maintain gains up to 7.4 years after treatment. These studies have not been replicated. In addition, little is known about the additional mental health care received by these patients in the intervening years after anxiety treatment, which seems likely to influence maintenance of gains for anxiety symptoms as well as the development of additional psychopathology during the subsequent years. The original sample consisted of 335 children ages 7 to 17 years (mean 13.09, 53% female) diagnosed with an anxiety disorder in 2010. Medical record review included provider billing records for mental health appointments during the five years after anxiety treatment. The subsample for this study was classified into three groups: 64 children who received CBT in an anxiety disorders clinic, 56 who received treatment from a psychiatrist, and 10 who were seen in a primary care setting. Chi-square analyses resulted in significant differences in mental health care utilization across the five years after treatment. Youth receiving treatment in primary care averaged less than one appointment each year and the appointments continued at the same rate across time. Children treated by a psychiatrist averaged approximately 3 appointments in the first two years and 2 in the subsequent three years. Importantly, youth treated in the anxiety clinic demonstrated a gradual decrease in mental health appointments across time. The nuanced differences will be presented in greater detail. The results of the current study have important implications for developing dissemination materials to help guide parents when they are selecting treatment for their children. By including all mental health appointments, this study recognizes that anxiety is often comorbid with additional diagnoses and that receiving evidence-based treatment may have long-term benefits that are associated with improvements in broader mental health. One important caveat might be that the acuity of mental health influenced the level of care sought by patients included in this study; however, taking this possibility into account, it seems those seeking care in a primary care setting continued to require similar care at the end of the study, indicating little improvement in symptoms was experienced.

Keywords : anxiety, children, mental health, outcomes

Conference Title : ICPS 2017 : International Conference on Psychology and Sociology

Conference Location : Vancouver, Canada

Conference Dates : August 07-08, 2017