

## Utilization of Complete Feed Based on Ammoniated Corn Waste on Bali Cattle Performance

**Authors :** Elihasridas, Rusmana Wijaya Setia Ninggrat

**Abstract :** This research aims to study the utilization of ammoniated corn waste complete ration for substitution basal ration of natural grass in Bali cattle. Four treatments (complete feed ration consisted of: R1=40% natural grass + 60% concentrate (control), R2= 50% natural grass+50% concentrate, R3=60% natural grass+40% concentrate and R4=40% ammoniated corn waste+60% concentrate) were employed in this experiment. This experiment was arranged in a latin square design. Observed variables included dry matter intake (DMI), average daily gain and feed conversion. Data were analyzed by using the Analysis of Variance following a 4 x 4 Latin Square Design. The DMI for R1 was 7,15kg/day which was significantly ( $P < 0,05$ ) higher than R2 (6,32 kg/day) and R3(6,07 kg/day), but was not significantly different ( $P < 0,05$ ) from R4 (7,01 kg/day). Average daily gain for R1(0,75 kg/day) which was significantly ( $P < 0,05$ ) higher than R2(0,66 kg/day) and R3 (0,61 kg/day),but was not significantly different ( $P > 0,05$ ) from R4(0,74 kg/day). Feed conversion was not significantly affected ( $P > 0,05$ ) by ration. It was concluded that ammoniated corn waste complete ration (40% ammoniated corn waste + 60% concentrate) could be utilized for substitution natural grass basal ration.

**Keywords :** ammoniated corn waste, bali cattle, complete feed, daily gain

**Conference Title :** ICRANMM 2017 : International Conference on Ruminant Animal Nutrition and Microbial Metabolism

**Conference Location :** Tokyo, Japan

**Conference Dates :** May 28-29, 2017