

Traumatic Spinal Cord Injury; Incidence, Prognosis and the Time-Course of Clinical Outcomes: A 12 Year Review from a Tertiary Hospital in Korea

Authors : Jeounghee Kim

Abstract : Objective: To describe the incidence of complication, according to the stage of Traumatic Spinal Cord Injury (TSCI) which was treated at Asan Medical Center (AMC), Korea. Hereafter, it should be developed in nursing management protocol of traumatic SCI. Methods. Retrospectively reviewed hospital records about the patients who were admitted AMC Patients with traumatic spinal cord injury until January 2005 and December 2016 were analyzed (n=97). AMC is a single institution of 2,700 beds where patients with trauma and severe trauma can be treated. Patients who were admitted to the emergency room due to spinal cord injury and who underwent intensive care unit, general ward, and rehabilitation ward. To identify long-term complications, we excluded patients who were operated on to other hospitals after surgery. Complications such as respiratory(pneumonia, atelectasis, pulmonary embolism, and others), cardiovascular (hypotension), urinary (autonomic dysreflexia, urinary tract infection (UTI), neurogenic bladder, and others), and skin systems (pressure ulcers) from the time of admission were examined through medical records and images. Results: SCI was graded according to ASIA scale. The initial grade was checked at admission. (grade A 55(56.7%), grade B 14(14.4)%, grade C 11(11.3%), grade D 15(15.5%), and grade E 2(2.1%). The grade was rechecked when the patient was discharged after treatment. (grade A 43(44.3%), grade B 15(15.5%), grade C 12(12.4%), grade D 21(21.6%), and grade E 6(6.2%). The most common complication after SCI was UTI 24cases (mean 36.5day), sore 24cases (40.5day), and Pneumonia which was 23 cases after 10days averagely. The other complications after SCI were neuropathic pain 19 cases, surgical site infection 4 cases. 53.6% of patient who had SCI were educated about intermittent catheterization at discharge from hospital. The mean hospital stay of all SCI patients was 61days. Conclusion: The Complications after traumatic SCI were developed at various stages from acute phase to chronic phase. Nurses need to understand fully the time-course of complication in traumatic SCI to provide evidence-based practice.

Keywords : spinal cord injury, complication, nursing, rehabilitation

Conference Title : ICNIC 2018 : International Conference on Nursing Information and Care

Conference Location : Zurich, Switzerland

Conference Dates : January 15-16, 2018