Utilization of Antenatal Care Services by Domestic Workers in Delhi

Authors : Meenakshi

Abstract : Background: The complications during pregnancy are the major cause of morbidity and deaths among women in the reproductive age group. Childbearing is the most important phase in women's lives that occur mainly in the adolescent and adult years. Maternal health, thus is an important issue as this as this is important phase is also productive time for women as they strive fulfill their capabilities as an individual, mothers, family members and also as a citizen. The objective of the study is to document the coverage of ANC and its determinants among domestic workers. Method: A survey of 300 domestic workers were carried in Delhi. Only respondents in the age group (15-49) and whose recent birth was of 5 years preceding the survey were included. Socio-demographic data and information on maternal health was collected from these respondents Information on ANC was collected from total 300 respondents. Standard of living index were composed based on households assists and similarly autonomy index was computed based on women decision making power in the households taking certain key variables. Cross tabulations were performed to obtain frequency and percentages. Potential socio-economic determinants of utilization of ANC among domestic workers were examined using binary logistic regressions. Results: Out of 300 domestic workers survey, only 70.7 per cent per cent received ANC. Domestic workers who married at age 18 years and above are 4 times more likely to utilize antenatal services during their last birth (***p< 0.01). Comparison to domestic workers with number of living children two or less, domestic workers with number of living children more than two are less likely to utilize antenatal care services (**p< 0.05). Domestic workers belonging to Other Backward Castes are more likely to utilize antenatal care services than domestic workers belonging to scheduled tribes ((**p< 0.05). Conclusion: The level of utilization of maternal health services are less among domestic workers is less, as they spend most of their time at the employers household. Though demonstration effect do have impact on their life styles but utilization of maternal health services is poor. Strategies and action are needed to improve the utilization of maternal health services among this section of workers as they are vulnerable because of no proper labour legislations.

Keywords : antenatal care, domestic workers, health services, maternal health, women's health **Conference Title :** ICMNCH 2017 : International Conference on Maternal, Newborn and Child Health **Conference Location :** Paris, France **Conference Dates :** November 20-21, 2017