

Incidence of Vulval, Vaginal and Cervical Disease in Rapid Access Clinic in a London Tertiary Hospital Setting

Authors : Kieren Wilson, Gulnaz Majeed

Abstract : NHS constitution gives rights to the patient with suspected cancer to be seen by a cancer specialist within 2 weeks of referral. Guys and St Thomas Hospital (GSTT) is one of the largest cancer centres in London. NICE guidelines have provided guidance for health professionals to refer patients appropriately to RAC. In GSTT suspected gynae cancer referrals are mostly by NHS e-Referral Service with some fax, emails as well as paper referrals. The objective of this study was to evaluate compliance with 2-week referral pathway with emphasis on one stop diagnostic service with supporting efficient pathways. A prospective evaluation over 3 months (1 Jan 2017 to 31 Mar 2017) was undertaken. There were 26 clinics, 761 patients were booked in the clinics with a DNA rate of 13% (n=101) hence 606 patients were seen. Majority of referrals were for post menopausal bleeding (PMB) 25% (n=194) followed by cervical, vaginal, vulval reasons 23% (n=179) (abnormal cytology excluded as patients directly referred to colposcopy unit in GSTT), ovarian 7% (n=54) and endometrial 5% (n=41). Women with new or previous established diagnosis of cancer were 24, cervical (n=17), vulva (n=6) and vagina (n=1). Multifocal preinvasive disease vulva (VIN), vagina (VAIN) and cervix (CIN) was confirmed in twenty-six patients 4% (high prevalence in HIV patients). Majority of cervical referrals: PCB (n=14), cervical erosion (n=7), polyps (n=9) and cervical cyst were benign. However, two women with PMB had cervical cancer. Only 2 out of 13 referrals with vaginal concerns had VAIN. One case with non-cervical glandular cytology was confirmed to have endometrial cancer. One stop service based on the diagnostic support of ultrasound, colposcopy and hysteroscopy was achieved in 54% (n=359). Patients were discharged to GP, benign gynaecology, endometriosis, combined vulval/dermatology clinic or gynae oncology. 33% (n=202) required a second visit, 12% (n=70) third visit, 3% (n=19) fourth visit, 1% (n=4) fifth visit and 1% (n=6) sixth visit. Main reasons for follow ups were the unavailability of diagnostic slots, patient choice, need for interpreters, the discussion following gynae MDM review for triage to benign gynae, delay in availability of diagnostic results like histology/MRI/CT. Recommendations following this study are multi disciplinary review of pathways with the availability of additional diagnostic procedure slots to aim for one stop service. Furthermore, establishment of virtual and telephone consultations to reduce follow ups.

Keywords : multifocal disease, post menopausal bleeding, preinvasive disease, rapid access clinic

Conference Title : ICCPC 2018 : International Conference on Cervical Pathology and Colposcopy

Conference Location : London, United Kingdom

Conference Dates : May 14-15, 2018