

Memory Types in Hemodialysis Patients: A Study Based on Hemodialysis Duration, Zahedan, South East of Iran

Authors : B. Sabayan, A. Alidadi, S. Ebrahimi, N. M. Bakhshani

Abstract : Neuropsychological problems are more common in hemodialysis (HD) patients than in healthy individuals. The aim of this study was to investigate the effect of long term HD on memory types of HD patients. To assess the different type of memory, we used memory parts of the Persian Papers and Pencil Cognitive assessment package (PCAP) and Addenbrooke's Cognitive Examination (ACE-R). Our study included 80 HD patients of whom 39 had less than six months of HD and 41 patients and another group which had a history of HD more than six months. The population had a mean age of 51.60 years old and 27.5% of them were female. The scores of patients who have been hemodialyzed for a long time (median time of HD was up to 4 years) had lower score in anterograde, explicit, visual, recall and recognition memory (5.44 ± 1.07 , 9.49 ± 3.472 , 22.805 ± 6.6913 , 5.59 ± 10.435 , 11.02 ± 3.190 score) than the HD patients who underwent HD for a shorter term, where the median time was 3 to 5 months ($P < 0.01$). The regression result shows that, by increasing the HD duration, all memory types are reduced ($R^2 = 0.600$, $P < 0.01$). The present study demonstrated that HD patients who were under HD for a long time had significantly lower scores in the different types of memory. However, additional researches are needed in this area.

Keywords : hemodialysis patients, duration of hemodialysis, memory types, Zahedan

Conference Title : ICNCS 2018 : International Conference on Neuropsychology and Cognitive Science

Conference Location : Tokyo, Japan

Conference Dates : May 28-29, 2018