

Rathke's Cleft Cyst Presenting as Unilateral Visual Field Defect

Authors : Ritesh Verma, Manisha Rath, Chand Singh Dhull, Sumit Sachdeva, Jitender Phogat

Abstract : A Rathke's cleft cyst is a benign growth found on the pituitary gland in the brain, specifically a fluid-filled cyst in the posterior portion of the anterior pituitary gland. It occurs when the Rathke's pouch does not develop properly and ranges in size from 2 to 40mm in diameter. A 38-year-old male presented to the outpatient department with loss of vision in the inferior quadrant of the left eye since 15 days. Visual acuity was 6/6 in the right eye and 6/9 in the left eye. Visual field analysis by HFA-24-2 revealed an inferior field defect extending to the supero-temporal quadrant in the left eye. MRI brain and orbit was advised to the patient and it revealed a well defined cystic pituitary adenoma indenting left optic nerve near optic chiasm consistent with the diagnosis of Rathke's cleft cyst (RCC). The patient was referred to neurosurgery department for further management. Symptoms vary greatly between individuals having RCCs. RCCs can be non-functioning, functioning, or both. Besides headaches, neurocognitive deficits are almost always present but have a high rate of immediate reversal if the cyst is properly treated or drained.

Keywords : pituitary tumors, rathke's cleft cyst, visual field defects, vision loss

Conference Title : ICOGT 2018 : International Conference on Ophthalmology and Glaucoma Treatments

Conference Location : Singapore, Singapore

Conference Dates : January 08-09, 2018