

## Antibacterial and Anti-Biofilm Activity of Papain Hydrolysed Camel Milk Whey and Its Fractions

**Authors :** M. Abdel-Hamid, P. Saporito, R. V. Mateiu, A. Osman, E. Romeih, H. Jenssen

**Abstract :** Camel milk whey (CMW) was hydrolyzed with papain from *Carica papaya* and fractionated by size exclusion chromatography (SEC). The antibacterial and anti-biofilm activity of the CMW, Camel milk whey hydrolysate (CMWH) and the obtained SEC-fractions was assessed against *Pseudomonas aeruginosa* and Methicillin-resistant *Staphylococcus aureus* (MRSA). SEC-F2 (fraction 2) exhibited antibacterial effectiveness against MRSA and *P. aeruginosa* with the minimum inhibitory concentration of 0.31 and 0.156 mg/ml, respectively. Furthermore, SEC-F2 significantly decreased biofilm biomass by 71% and 83 % for MRSA and *P. aeruginosa* in a crystal violet microplate assay. Scanning electron microscopy showed that the SEC-F2 caused changes in the treated bacterial cells. Additionally, LC/MS analysis was used to characterize the peptides of SEC-F2. Two major peptides were detected in SEC-F2 having masses of 414.05 Da and 456.06 Da. In conclusion, this study has demonstrated that hydrolysis of CMW with papain generates small and extremely potent antibacterial and anti-biofilm peptides against both MRSA and *P. aeruginosa*.

**Keywords :** camel milk, whey proteins, antibacterial peptide, anti-biofilm

**Conference Title :** ICFMFS 2018 : International Conference on Food Microbiology and Food Safety

**Conference Location :** Prague, Czechia

**Conference Dates :** March 22-23, 2018