Predatory Potential of Coccinella septempunctata Linnaeus and Coccinella undecimpunctata Linnaeus on Different Prey Species

Authors : Adnan A. E. Darwish

Abstract : The predatory potential and preference of both larvae and adult of seven-spot ladybird, Coccinella septempunctata Linnaeus and the eleven-spot ladybird, Coccinella undecimpunctata Linnaeus to the green peach aphid, Myzus persicae (Sulzer), the cotton aphid, Aphis gossypii Glover, the bird cherry-oat aphid, Rhopalosiphum padi (Linnaeus) and onion thrips, Thrips tabaci Lindeman were investigated under laboratory conditions at varying prey densities at faculty of Agriculture, Damanhour university, Egypt. There were significant differences between the consumed numbers of the four different species by the two different lady beetle species. The most consumed prey by C. septempunctata was the A. gossypii followed by R. padi then M. persicae and finally T. tabaci and these results were repeated in case of C. undecimpunctata. As the grubs of C. septempunctata and C. undecimpunctata developed from 1st to 4th larval instars, the consumption rate from aphid species and thrips increased. The consumption rate of M. persicae, A. gossypii, R. padi and T. tabaci significantly increased with the advancement in the larval stage of the predator. The forth larval instar of C. septempunctata and C. undecimpunctata exhibited the highest predatory potential comparing to the first, second and third larval instars. The number of prey eaten by adult stage or different instars of larvae of the two predators increased significantly with prey density, reaching the maximum value when 150 preys were provided compared with 50 and 100 preys.

Keywords : predatory potential, Coccinella septempunctata, Coccinella undecimpunctata, Thrips tabaci, Myzus persicae, Aphis gossypii, Rhopalosiphum padi

Conference Title : ICE 2018 : International Conference on Entomology

Conference Location : Paris, France

Conference Dates : October 29-30, 2018