

Inactivation of Root-Knot Nematode Eggs *Meloidogyne enterolobii* in Irrigation Water Treated with Ozone

Authors : I. A. Landa-Fernandez, I. Monje-Ramirez, M. T. Orta-Ledesma

Abstract : Every year plant-parasitic nematodes diminish the yield of high-value crops worldwide causing important economic losses. Currently, *Meloidogyne enterolobii* has increased its importance due to its high aggressiveness, increasing geographical distribution and host range. Root-knot nematodes inhabit the rhizosphere soil around plant roots. However, they can come into contact with irrigation water. Thus, plant-parasitic nematodes can be transported by water, as eggs or juveniles. Due to their high resistance, common water disinfection methods are not effective for inactivating these parasites. Ozone is the most effective disinfectant for microbial inactivation. The objective of this study is to demonstrate that ozone treatment is an alternative method control in irrigation water of the root-knot nematode *M. enterolobii*. It has been shown that ozonation is an effective treatment for the inactivation of protozoan cysts and oocysts (*Giardia* and *Cryptosporidium*) and for other species of the genus *Meloidogyne* (*M. incognita*), but not for the *enterolobii* specie. In this study, the strain of *M. enterolobii* was isolated from tomatoes roots. For the tests, eggs were used and were inoculated in water with similar characteristics of irrigation water. Subsequently, the disinfection process was carried out in an ozonation unit. The performance of the treatments was evaluated through the egg's viability by assessing its structure by optical microscopy. As a result of exposure to ozone, the viability of the nematode eggs was reduced practically in its entirety; with dissolved ozone levels in water close to the standard concentration (equal to 0.4 mgO₃/L), but with high contact times (greater than 4 min): 0.2 mgO₃/L for 15 minutes or 0.55 mgO₃/L for 10 minutes. Additionally, the effect of temperature, alkalinity and organic matter of the water was evaluated. Ozonation is effective and a promising alternative for the inactivation of nematodes in irrigation water, which could contribute to diminish the agricultural losses caused by these organisms.

Keywords : inactivation process, irrigation water treatment, ozonation, plant-parasite nematodes

Conference Title : ICEWW 2018 : International Conference on Environment, Water and Wetlands

Conference Location : Toronto, Canada

Conference Dates : June 21-22, 2018