

Effectiveness of Control Measures for Ambient Fine Particulate Matters Concentration Improvement in Taiwan

Authors : Jiun-Horng Tsai, Shi-Jie, Nieh

Abstract : Fine particulate matter (PM_{2.5}) has become an important issue all over the world over the last decade. Annual mean PM_{2.5} concentration has been over the ambient air quality standard of PM_{2.5} (annual average concentration as 15µg/m³) which adapted by Taiwan Environmental Protection Administration (TEPA). TEPA, therefore, has developed a number of air pollution control measures to improve the ambient concentration by reducing the emissions of primary fine particulate matter and the precursors of secondary PM_{2.5}. This study investigated the potential improvement of ambient PM_{2.5} concentration by the TEPA program and the other scenario for further emission reduction on various sources. Four scenarios had been evaluated in this study, including a basic case and three reduction scenarios (A to C). The ambient PM_{2.5} concentration was evaluated by Community Multi-scale Air Quality modelling system (CMAQ) ver. 4.7.1 along with the Weather Research and Forecasting Model (WRF) ver. 3.4.1. The grid resolutions in the modelling work are 81 km × 81 km for domain 1 (covers East Asia), 27 km × 27 km for domain 2 (covers Southeast China and Taiwan), and 9 km × 9 km for domain 3 (covers Taiwan). The result of PM_{2.5} concentration simulation in different regions of Taiwan shows that the annual average concentration of basic case is 24.9 µg/m³, and are 22.6, 18.8, and 11.3 µg/m³, respectively, for scenarios A to C. The annual average concentration of PM_{2.5} would be reduced by 9-55 % for those control scenarios. The result of scenario C (the emissions of precursors reduce to allowance levels) could improve effectively the airborne PM_{2.5} concentration to attain the air quality standard. According to the results of unit precursor reduction contribution, the allowance emissions of PM_{2.5}, SO_x, and NO_x are 16.8, 39, and 62 thousand tons per year, respectively. In the Kao-Ping air basin, the priority for reducing precursor emissions is PM_{2.5} > NO_x > SO_x, whereas the priority for reducing precursor emissions is PM_{2.5} > SO_x > NO_x in others area. The result indicates that the target pollutants that need to be reduced in different air basin are different, and the control measures need to be adapted to local conditions.

Keywords : airborne PM_{2.5}, community multi-scale air quality modelling system, control measures, weather research and forecasting model

Conference Title : ICAPC 2019 : International Conference on Air Pollution and Control

Conference Location : London, United Kingdom

Conference Dates : May 23-24, 2019